

# Distribution of CC-chemokine receptor-5- $\Delta$ 32 allele among the tribal and caste population of Vidarbha region of Maharashtra state

Arvind B. Chavhan<sup>1,2</sup>, Santosh S. Pawar<sup>3</sup>, Rajusing G. Jadhao<sup>1</sup>, Kishor G. Patil<sup>2</sup>

Departments of Zoology, <sup>1</sup>Shri Shivaji Science College, Amravati, <sup>2</sup>Institute of Science, Nagpur, <sup>3</sup>Govt. Vidarbha Institute of Science and Humanities, Amravati, Maharashtra, India

**BACKGROUND:** Genetic relationships among the ethnic groups are not uniform across the geographical region. Considering this assumption, we analyzed the frequency of the CC-chemokine receptor-5 (CCR5)- $\Delta$ 32 allele of the CCR5 chemokine receptor, which is considered a Caucasian marker, in *Bhil* tribal and *Brahmin* caste sample sets from the population.

**MATERIALS AND METHODS:** 108 blood samples were collected from 6 tribe's populations and a caste population from the district of Vidarbha region.

**RESULTS AND DISCUSSION:** The presence of low frequencies of CCR5- $\Delta$ 32 in an individual of *Bhil* tribe (0.034,  $\chi^2$  value 0.017) in the present study implies that these communities may have a better resistance toward human immunodeficiency virus (HIV)/acquired immunodeficiency syndrome (AIDS) than the other studied tribe sample, as non-show such mutation.

**CONCLUSION:** The marginal presence of the allele seen in the studied tribal population could be due to gene flow from the people of European descent. However, lack of the homozygous CCR5- $\Delta$ 32 mutation and the low prevalence of heterozygous CCR5- $\Delta$ 32 mutations suggest that the Indians are highly susceptible to HIV/AIDS, and this correlates with the highest number of HIV/AIDS infected individuals in India.

**Key words:** Allele frequency, CC-chemokine receptor-5- $\Delta$ 32, India, genetic polymorphism, tribes, Vidarbha

## Introduction

Human immunodeficiency virus (HIV)-1 infection has spread to all population groups in India and has reached epidemic proportions.<sup>[1]</sup> The rate of progression of HIV-1 disease exhibits a remarkable variation among different individuals. Many host genetic factors are now known to affect the disease progression rates, especially polymorphisms in genes encoding chemokine receptors.<sup>[2-5]</sup> Although, no studies on chemokine receptor polymorphisms have been reported in the endogamous population of State of Maharashtra state so far, only one study has been carried out in healthy individuals of tribes and Muslim ethnic groups of Andhra Pradesh, south India.<sup>[6]</sup>

Certain members of the chemokine family of receptors serve as critical portals for the entry of HIV-1 into target cells. A mutant allele (CC-chemokine receptor-5 [CCR5]- $\Delta$ 32) of the  $\beta$ -chemokine receptor gene CCR5 carrying a 32 base-pair deletion prevents cell invasion by the primary transmitting strain of HIV-1. Individuals who are homozygous for the CCR5- $\Delta$ 32/ $\Delta$ 32 allele are highly resistant to HIV-1 infection; the heterozygote state does not protect against HIV-1 infection, although, heterozygotes have been found to have significantly lower viral loads.<sup>[7]</sup> Early reports indicated that the CCR5- $\Delta$ 32 allele maybe absent in indigenous non-European populations. The CCR5\_32 allele appears to have originated quite recently (approximately 7,000 years ago) in northeastern

### Access this article online

Quick Response Code:



Website:

www.ijhg.com

DOI:

10.4103/0971-6866.112894

**Address for correspondence:** Prof. A. B. Chavhan, Department of Zoology, Shri Shivaji Science College, Morshi Road, Amravati, Maharashtra, India. E-mail: arvind.bioinfo26@gmail.com

Europe.<sup>[8]</sup> Although its frequency has now reached a relatively high level in Europeans, e.g., 16.3% in Finns and 15.8% in Moravians, it is not present among African populations, and is only so at low levels in the Asian.<sup>[9]</sup> Hence, by application it is possible to evaluate the influence of the European population on the genetic constitution of others.

In India population one study has been carried out in healthy individuals of the tribes and Muslim ethnic groups of Andhra Pradesh, South India.<sup>[6]</sup> Majumder and Dey,<sup>[10]</sup> reported absence of CCR5-Δ32 in various ethnic populations of India, both tribal and non-tribal, except for some populations of the northern and western regions where this allele may have been introduced by Caucasian gene flow. Although a few studies on chemokine, chemokine receptor, and DCSIGN exon 4 repeat number polymorphisms have been reported in north Indian (Aryan descent) HIV patients and healthy controls,<sup>[11-13]</sup> there is a dearth of reports on the HIV patients with and without tuberculosis (TB) of South Indian (Dravidian descent) origin.

CCR5-Δ32 exhibit variable frequencies in distinct populations<sup>[14-17]</sup> and possibly, their phenotypes depend on the ethnicity analyzed.<sup>[15-20]</sup> The Brazilian population presents a complex genetic background, characterized by a high degree of miscegenation.<sup>[21,22]</sup> In major Brazilian cities, CCR5-Δ32 was found at frequencies of between 2% and 7%.<sup>[23-26]</sup>

Genetic relationships among caste groups are not uniform across the geographical region of India.<sup>[27,28]</sup> India is known for the enormous cultural and genetic diversity of its people.<sup>[27]</sup> Such diversity is some time attributed to the positioning of the Indian peninsula at the tri-junction of the three continents, viz. Africa, Europe, and Asia. The contemporary Indian population is stratified as tribal and non-tribal, i.e., caste population. The origin of the caste in India is an enigma,<sup>[29]</sup> though many are known to have a tribal origin.<sup>[30,31]</sup>

The Maharashtra state of India forms a huge irregular triangle with its base on the west coast of India, overlooking the Arabian Sea. Historically, the state is comprised of three sub-regions, Western Maharashtra, the Vidarbha, and the Marathwada. Vidarbha lies on the eastern side and thus mainly contributes to the region

broadly referred to as central India. Apart from the tribal population, many other Ethnic Communities mainly Hindus, Muslim, Buddhist and Sikhs, inhabit the region. The Vidarbha has a hoary past and has been under the domination of many Hindus, Muslims, and tribal-Gond Kingdoms. The Vidarbhan strip served as a bridge between Northern and Southern India. It is assumed that the relationship between these various populations may define the present genetic landscape of India.

Taking this assumption and geographical and ethnic diversity into account in the present study, we investigated the distribution of CCR5-Δ32 alleles in tribes from the Vidarbha region (Maharashtra) Central India.

## Material and Methods

### Population

#### The Kolam (tribes)

Besides inhabiting the adjoining state, a substantial number of these people inhabit in few district of Vidarbha. They speak the Gondi dialect which belongs to the Dravidian linguistic group. We sampled 15 samples from this group Village of Yavatmal district.

#### The Bhil (tribes)

*Bhils* are listed as Adivasi residents of the states of Gujarat, Madhya Pradesh, Chhattisgarh, Maharashtra, and Rajasthan in Western and Central India as well as in Tripura in far-eastern India on the border with Bangladesh. *Bhils* are divided into a number of endogamous territorial divisions, which in turn have a number of clans and lineages. Most *Bhils* now speak the language of the region they reside in, such as Marathi and Gujarati. We sampled 15 samples from this group Village of Yavatmal district.

#### The Korkus (tribes)

The *Korkus* are a typical tribal population from Amravati district and found only in the *Satpuda* mountain ranges spanning Maharashtra and Madhya Pradesh. They are mainly concentrated in *Melghat* a scheduled area of *Korku* comprising 89% of the tribal population. The *Korkus* speak *Korku* dialect belonging to Austro-Asiatic linguistic group. The Austro-Asiatic speakers are considered as the first settler of Indian subcontinent.<sup>[32]</sup> We sampled 15 samples from this group.

### The *Paradhi* (tribes)

Phase *Paradhi* or Phasse *Paradhi* is a tribe in India. The tribe often faces harassment by Indian law enforcement agencies. The tribe is found mostly in Maharashtra and parts of Madhya Pradesh. The Phasse are a sub tribe of the *Paradhi* caste, which includes sub-castes like *Gav Paradhi*, *Berad-Paradhi*, *Gay-Paradhi*, *Chita Paradhi*. *Paradhi* is the term for “hunter.” There are only three surnames among them, Chauhan, Pawar, and Solanke. We sampled 15 samples from this group from Akola district.

### The *Andh* (tribes)

A low cultivating caste of Berar, who numbered 52,000 persons in 1911, and belongs to the Yeotmal, Akola, and Buldana Districts. The *Andhs* appear to be a non-Aryan tribe of the Andhra or Tamil country, from which they derive their name. There were 8228 *Andh* in Andhra Pradesh in 1991. According to Singh *et al.*,<sup>[33]</sup> in their 2004 book people of India there are over 74,000 *Andhs* in Maharashtra. We sampled 16 samples from this group.

### The *Gonds* (tribes)

This tribe falls under the primitive tribes category and spread much over the central India. *Gond* generally speaks “*Gondi*” dialect, which belongs to the Dravidian linguistic family, after Indo-European, in India. We sampled 17 blood samples from this tribe from the Village Gadchiroli district.

### *Brahmin* caste

*Brahmin* is a class of priests and preachers of ‘*Dharma*’ and considered as the torch bearer of Hinduism. Majority of *Brahmin* in Maharashtra speak Marathi, one of the major languages of Indo-Aryan linguistic group. The population of *Brahmins* in Amravati district is 21,500 or 3% of the population. We collected 15 samples of *Brahmins* living in Amravati district.

### Blood sample collection and DNA extraction

We have collected 108 blood samples on Whatman FTA mini cards (GE Health-Care, UK, Ltd) from 6 tribe’s populations and a caste population from the district of

Vidarbha region. Every card was labeled with appropriate code as per tribes/caste and district with informed consent obtained from each volunteer. Approximately, (200 µl [2-5 drop]) of blood by veni-puncture was directly spotted on the FTA mini card within a printed circle area and dried at room temperature. Genomic DNA was isolated from each dried blood sample by following protocol.<sup>[34]</sup>

### Amplification and restriction digestion of DNA

Polymerase chain reaction was performed following previously prescribed protocol by.<sup>[35]</sup> Briefly, 100 ng of genomic DNA was denatured at 94°C for 10 min, following, which it was subjected to 30 cycles of denaturation, annealing, and extension. The last cycle was followed by an incubation at 72°C for 10 min. The reaction mixture of 50 µl contained, 50 mmol KCl, 10 mmol Tris-HCl, pH 8.3, 800 µmol dNTPs, 100 µg/ml gelatin, 10 pmoles of each of the CCR5-specific primer forward: CCR5 – F: (5'-CCTGGCTGTCGTCATGCTG-3') and reverse: CCR5-R: (5'-CTGATCTAGAGCCATGTGCAC AACTCT-3')., and 1.5 units of Taq polymerase enzyme (Xcelris Genomics, Ahmedabad, India).

### Genotyping for CCR5 Δ-32 polymorphism

The genotypes were visualized by running digested product on 2% agarose gel at 100 V for about 2 h and the results were recorded in gel documentation system. The *EcoRI* Restriction enzymes digest the amplified polymerization Chain Reaction (PCR) product of 735 base pairs (bps). The amplified product was digested with 10 U of *EcoRI* at 37°C for 2 h. After digestion, the products were analyzed on a 2% agarose gel and bands were visualized on a Ultraviolet (UV)-transilluminator. PCR amplified a 735 bp region of genomic DNA that spanned a 32-bp deletion differentiating the CCR5-<sub>32</sub> allele from its wild-type counterparts at the CCR5 locus. After restriction digestion with *EcoRI*, with wild gene yielded band at 332 bp and for mutated gene, the bands were at 332 and 403 Figure 1.

### Statistical analysis

Statistical analysis of allele frequencies was performed using Chi-square statistics. Genotype distribution for polymorphism was first compared to predictable values

from Hardy–Weinberg equilibrium. In all cases, *P* values less than 0.05 were considered to be statistically significant.

### Observation and Results

Our data on distribution of CCR5-Δ32 mutation among the selected tribe and a caste is depicted Table 1. Genotype and phenotype for the heterozygous mutation among the tribes sample suggested that it is either absent or present at low frequency 1.08% (1 in 93 tribe's samples and 1 in 15 samples of a caste). None of the tribes and control caste was found to be homozygous for the CCR5-Δ32 mutation, while the *Bhil* tribe and control caste show the heterozygous for the CCR5-Δ32 mutation in negligible frequency (0.034). The analysis of  $\chi^2$  suggested that the prevalence of the CCR5-Δ32 is significantly low ( $\chi^2 = 0.02$ ,  $P > 0.05$ ). The aggregate frequencies of the entire sample for the wild-type allele CCR5 and the CCR5-Δ32 variant were found to be 0.991 and 0.009, respectively. Among the tribes *Kolam*, *Korku*,


*Paradhi*, *Andh* and *Gonds* showed the highest (wild type) homozygous genotype frequency (100%). while only the *Bhil* tribe shows heterozygous genotype frequency (06.97%) and allelic frequency (0.034) for CCR5/Δ32. Only one bearer of this mutation was found in the blood sample of *Bhil* tribe collected from Yavatmal district of Vidarbha region.

No significant deviations from the HWE were observed ( $P > 0.05$ , Chi-squared goodness of fit). Table 1 shows the frequency of the CCR5-Δ32 allele in the six tribes and one control caste *Brahmin* population.

### Discussion on Conclusion

This study describes the genotype and allele frequencies of the polymorphisms CCR5-Δ32 in selected six tribal and one caste populations from Vidarbha region. Most importantly, however, this study is the first to be conducted in Vidarbha that investigates the genetic polymorphisms CCR5-Δ32 among different ethnic tribal population settlement in the districts of Vidarbha region.

CCR5, a coreceptor for HIV-I virus, has been shown to be the most important for the HIV transmission.<sup>[36,37]</sup> A 32-nucleotide deletion of CCR5 homozygous (CCR5-Δ32/Δ32) display a high degree of the natural resistance to HIV transmission whereas CCR5-Δ32 heterozygosity (CCR5+/Δ32) demonstrate a slower progression to acquired immunodeficiency syndrome (AIDS) than CCR5 wild type (CCR5+/+).<sup>[3,38]</sup> However, this genetic mutation is found in Caucasian rather than non-Caucasian population including India.<sup>[10]</sup> The CCR5-Δ32 genotype frequency among our study tribes sample was absent or negligible, the average genotype frequency were common homozygous



**Figure 1: CC-chemokine receptor-5 genotyping among the tribes Lane 1, 2, 3, 5, 6 and 7 represent the PCR product from samples with homozygous wild type genotypes (fragments of 332 bp wt/wt). Lane 4 represents the Δ-32 genotype (with the presence of both fragment of 332 and 403 wt/Δ32)**

**Table 1: Genotypic distribution and gene frequencies of the CCR5 allele in different population samples of Vidarbha region of Maharashtra state**

Population	Genotype of CCR5 Δ32			Allelic frequency ( $\lambda$ )		Δ32	$\chi^2$	<i>P</i> value (1 df)
	wt/wt	wt/Δ32	Δ32/Δ32	<i>P</i> <sup>2</sup>	<i>q</i> <sup>2</sup>			
<i>Kolam</i>	15 (100)	0	0	1.00	0.00	0	NA	NA
<i>Bhil</i>	14 (93.3)	1 (06.9)	0	0.966	0.034	0.034	0.017	0.893
<i>Korku</i>	15 (100)	0	0	1.00	0.00	0	NA	NA
<i>Paradhi</i>	15 (100)	0	0	1.00	0.00	0	NA	NA
<i>Andh</i>	16 (100)	0	0	1.00	0.00	0	NA	NA
<i>Gond</i>	17 (100)	0	0	1.00	0.00	0	NA	NA
<i>Brahmin</i>	14 (93.3)	1 (06.9)	0	0.966	0.034	0.034	0.02	0.893
Total	106 (98.1)	2 (01.8)	0	0.991	0.009	0.01	0.009	0.922

NA: Not applicable, CCR5: CC-chemokine receptor-5

wt/wt (98.3%), heterozygous wt/mt (1.87%) and rare homozygous mt/mt (0%), from the control group, revealing CCR5-Δ32 allele frequency of 6.97%. The frequency of the CCR5-Δ32 allele among our study population seems to be remarkably similar to previously reported frequencies in other Asian populations.

The CCR5-Δ32 allele frequency among Asians is very low in Rajasthan Indians (0.05%), Andhra Pradesh Indians (0-0.03%),<sup>[39]</sup> North Indians (1.5%),<sup>[12]</sup> South Indians (1-3%),<sup>[6]</sup> and ethnic population of Kashmir (3-4%).<sup>[40]</sup> A similar study conducted from the Island of Crete, Greece showed allele frequency of 3.25%, with a 95% confidence interval (CI) for conformity with Hardy-Weinberg equilibrium of 0.74-5.7%.<sup>[41]</sup> The CCR5-Δ32 polymorphism is found all across Europe at different allele frequencies, with a North to South decreasing gradient and lower distribution in the regions of Southeast Mediterranean.<sup>[42]</sup> The frequency of the CCR5-Δ32 allele in the studied tribal population is consistently similar with data reported from other populations with non-European ancestors.<sup>[9,16]</sup> CCR5-Δ32 allelic frequencies were not different when the self-reported racial characteristics of the individuals evaluated were considered.<sup>[25]</sup> This allele has not been found among South American native Indians, corroborating the hypothesis of a European origin of this allele and its introduction to the continent through migration.<sup>[43]</sup>

Within the Middle-Eastern populations the frequency of the mutant CCR5-Δ32 allele reached it's the highest among Iranians, 2.4%; Saudi, 2.1%; and it's the lowest among Kuwaitis, 1%; and the Egyptians, 0.5%; and is completely absent in individuals from the United Arab Emirates.<sup>[44]</sup> Our results suggest that the CCR5-Δ32 allele is detected at very low frequency in studied tribal populations from Vidarbha. The presence of low frequencies of CCR5-Δ32 in an individual of *Bhil* tribe (0.034,  $\chi^2$  value 0.017) in the present study implies that these communities may have a better resistance to HIV/AIDS than other studied tribe sample, as non show such mutation. However, the CCR5-Δ32 allele is observed mostly in European populations.

The marginal presence of the allele seen in the studied tribal population could be due to gene flow from the people of European descent. However, CCR5-Δ32

is completely absent in the populations from Africa, Oceania, and the Americas. However, lack of the homozygous CCR5-Δ32 mutation and the low prevalence of heterozygous CCR5-Δ32 mutations suggest that the Indians are highly susceptible to HIV/AIDS, and this correlates with the highest number of HIV/AIDS infected individuals in India.

## References

- Misra SN, Sengupta D, Satpathy SK. AIDS in India: Recent trends in opportunistic infections. *Southeast Asian J Trop Med Public Health* 1998;29:373-6.
- Berger EA, Murphy PM, Farber JM. Chemokine receptors as HIV-1 coreceptors: Roles in viral entry, tropism, and disease. *Annu Rev Immunol* 1999;17:657-700.
- O'Brien TR, Goedert JJ. Chemokine receptors and genetic variability: Another leap in HIV research. *JAMA* 1998;279:317-8.
- Moore JP. Coreceptors: Implications for HIV pathogenesis and therapy. *Science* 1997;276:51-2.
- Fauci AS. Host factors and the pathogenesis of HIV-induced disease. *Nature* 1996;384:529-34.
- Ramana GV, Vasanthi A, Khaja M, Su B, Govindaiah V, Jin L, et al. Distribution of HIV-1 resistance-conferring polymorphic alleles SDF-1-3'A, CCR2-64I and CCR5-Delta32 in diverse populations of Andhra Pradesh, South India. *J Genet* 2001;80:137-40.
- Huang Y, Paxton WA, Wolinsky SM, Neumann AU, Zhang L, He T, et al. The role of a mutant CCR5 allele in HIV-1 transmission and disease progression. *Nat Med* 1996;2:1240-3.
- Lidén K, Linderholm A, Götherström A. Pushing it back. Dating the CCR5-D32 bp deletion to the mesolithic in Sweden and its implications for the Meso/Neo transition. *Doc Praehist* 2006;33:29-37.
- Martinson JJ, Chapman NH, Rees DC, Liu YT, Clegg JB. Global distribution of the CCR5 gene 32-basepair deletion. *Nat Genet* 1997;16:100-3.
- Majumder PP, Dey B. Absence of the HIV-1 protective Delta ccr5 allele in most ethnic populations of India. *Eur J Hum Genet* 2001;9:794-6.
- Suresh P, Wanchu A, Sachdeva RK, Bhatnagar A. Gene polymorphisms in CCR5, CCR2, CX3CR1, SDF-1 and RANTES in exposed but uninfected partners of HIV-1 infected individuals in North India. *J Clin Immunol* 2006;26:476-84.
- Verma R, Gupta RB, Singh K, Bhasin R, Anand Shukla A, Chauhan SS, et al. Distribution of CCR5delta32, CCR2-64I and SDF1-3'A and plasma levels of SDF-1 in HIV-1 seronegative North Indians. *J Clin Virol* 2007;38:198-203.
- Rathore A, Chatterjee A, Sivarama P, Yamamoto N, Singhal PK, Dhole TN. Association of RANTES-403 G/A,-28 C/G and In1.1 T/C polymorphism with HIV-1 transmission and progression among North Indians. *J Med Virol* 2008;80:1133-41.
- Salem AH, Farid E, Fadel R, Abu-Hijleh M, Almawi W, Han K, et al. Distribution of four HIV type 1-resistance

- polymorphisms (CCR5-Delta32, CCR5-m303, CCR2-64I, and SDF1-3'A) in the Bahraini population. *AIDS Res Hum Retroviruses* 2009;25:973-7.
15. Gonzalez E, Dhanda R, Bamshad M, Mummidi S, Geevarghese R, Catano G, *et al.* Global survey of genetic variation in CCR5, RANTES, and MIP-1alpha: Impact on the epidemiology of the HIV-1 pandemic. *Proc Natl Acad Sci U S A* 2001;98:5199-204.
  16. Su B, Sun G, Lu D, Xiao J, Hu F, Chakraborty R, *et al.* Distribution of three HIV-1 resistance-conferring polymorphisms (SDF1-3'A, CCR2-64I, and CCR5-delta32) in global populations. *Eur J Hum Genet* 2000;8:975-9.
  17. Smith MW, Dean M, Carrington M, Winkler C, Huttley GA, Lomb DA, *et al.* Contrasting genetic influence of CCR2 and CCR5 variants on HIV-1 infection and disease progression. Hemophilia Growth and Development Study (HGDS), Multicenter AIDS Cohort Study (MACS), Multicenter Hemophilia Cohort Study (MHCS), San Francisco City Cohort (SFCC), ALIVE Study. *Science* 1997;277:959-65.
  18. Gonzalez E, Bamshad M, Sato N, Mummidi S, Dhanda R, Catano G, *et al.* Race-specific HIV-1 disease-modifying effects associated with CCR5 haplotypes. *Proc Natl Acad Sci U S A* 1999;96:12004-9.
  19. Mummidi S, Ahuja SS, Gonzalez E, Anderson SA, Santiago EN, Stephan KT, *et al.* Genealogy of the CCR5 locus and chemokine system gene variants associated with altered rates of HIV-1 disease progression. *Nat Med* 1998;4:786-93.
  20. An P, Martin MP, Nelson GW, Carrington M, Smith MW, Gong K, *et al.* Influence of CCR5 promoter haplotypes on AIDS progression in African-Americans. *AIDS* 2000;14:2117-22.
  21. Callegari-Jacques SM, Grattapaglia D, Salzano FM, Salamoni SP, Crossetti SG, Ferreira ME, *et al.* Historical genetics: Spatiotemporal analysis of the formation of the Brazilian population. *Am J Hum Biol* 2003;15:824-34.
  22. Pena SD. Reasons for banishing the concept of race from Brazilian medicine. *Hist Cienc Saude Manguinhos* 2005;12:321-46.
  23. Munerato P, Azevedo ML, Sucupira MC, Pardini R, Pinto GH, Catroxo M, *et al.* Frequency of polymorphisms of genes coding for HIV-1 co-receptors CCR5 and CCR2 in a Brazilian population. *Braz J Infect Dis* 2003;7:236-40.
  24. Vargas AE, Marrero AR, Salzano FM, Bortolini MC, Chies JA. Frequency of CCR5delta32 in Brazilian populations. *Braz J Med Biol Res* 2006;39:321-5.
  25. Reiche EM, Ehara Watanabe MA, Bonametti AM, Morimoto HK, Akira Morimoto A, Wiechmann SL, *et al.* Frequency of CCR5-Delta32 deletion in human immunodeficiency virus type 1 (HIV-1) in healthy blood donors, HIV-1-exposed seronegative and HIV-1-seropositive individuals of southern Brazilian population. *Int J Mol Med* 2008;22:669-75.
  26. Rigato PO, Hong MA, Casseb J, Ueda M, de Castro I, Benard G, *et al.* Better CD4+ T cell recovery in Brazilian HIV-infected individuals under HAART due to cumulative carriage of SDF-1-3'A, CCR2-V64I, CCR5-D32 and CCR5-promoter 59029A/G polymorphisms. *Curr HIV Res* 2008;6:466-73.
  27. Majumder PP. People of India: Biological diversity and affinities. *Evol Anthropol* 1998;6:100-10.
  28. Baig MM, Khan AA, Kulkarni KM. Mitochondrial DNA diversity in tribal and caste groups of Maharashtra (India) and its implication on their genetic origins. *Ann Hum Genet* 2004;68:453-60.
  29. Majumder PP. Indian caste origins: Genomic insights and future outlook. *Genome Res* 2001;11:931-2.
  30. Karve I. Hindu Society-An Interpretation. Poona: Deshmukh Prakashan; 1961.
  31. Kasambi DD. The Culture and Civilization of Ancient India in Historical Outline. New Delhi: Vikas; 1964.
  32. Roychoudhury S, Roy S, Basu A, Banerjee R, Vishwanathan H, Usha Rani MV, *et al.* Genomic structures and population histories of linguistically distinct tribal groups of India. *Hum Genet* 2001;109:339-50.
  33. Singh S, Bhanu B. Anthropological Survey of India. Maharashtra: Popular Prakashan; 2004. p. 65.
  34. Zhou H, Hickford JG, Fang Q. A two-step procedure for extracting genomic DNA from dried blood spots on filter paper for polymerase chain reaction amplification. *Anal Biochem* 2006;354:159-61.
  35. Bernard NF, Yannakis CM, Lee JS, Tsoukas CM. Human immunodeficiency virus (HIV)-specific cytotoxic T lymphocyte activity in HIV-exposed seronegative persons. *J Infect Dis* 1999;179:538-47.
  36. Deng H, Liu R, Ellmeier W, Choe S, Unutmaz D, Burkhardt M, *et al.* Identification of a major co-receptor for primary isolates of HIV-1. *Nature* 1996;381:661-6.
  37. Dragic T, Trkola A, Lin SW, Nagashima KA, Kajumo F, Zhao L, *et al.* Amino-terminal substitutions in the CCR5 coreceptor impair gp120 binding and human immunodeficiency virus type 1 entry. *J Virol* 1998;72:279-85.
  38. Eugen-Olsen J, Iversen AK, Garred P, Koppelhus U, Pedersen C, Benfield TL, *et al.* Heterozygosity for a deletion in the CKR-5 gene leads to prolonged AIDS-free survival and slower CD4 T-cell decline in a cohort of HIV-seropositive individuals. *AIDS* 1997;11:305-10.
  39. Kozhekbaeva GM, Borodina TA, Borinskaia SA, Gusar VA, Feshchenko SP, Akhmetova VL, *et al.* Distribution of the HIV-1 resistance-conferring alleles (CCR5delta32, CCR2-64I, and SDF1 3'A) in Russian, Ukrainian, and Belarusian populations. *Genetika* 2004;40:1394-401.
  40. Irtiza S, Dil-Afroze, Naykoo NA, Lateef C, Iqbal Q, Inayat SF, *et al.* Polymorphism in the CC-chemokine receptor-5 (CCR5) gene and risk of AIDS among Kashmiri population. *J AIDS HIV Res* 2011;3:103-6.
  41. Apostolakis S, Baritaki S, Krambovitis E, Spandidos DA. Distribution of HIV/AIDS protective SDF1, CCR5 and CCR2 gene variants within Cretan population. *J Clin Virol* 2005;34:310-4.
  42. Libert F, Cochaux P, Beckman G, Samson M, Aksenova M, Cao A, *et al.* The deltaccr5 mutation conferring protection against HIV-1 in Caucasian populations has a single and recent origin in Northeastern Europe. *Hum Mol Genet* 1998;7:399-406.
  43. Leboutte AP, de Carvalho MW, Simões AL. Absence of the deltaccr5 mutation in indigenous populations of the Brazilian Amazon. *Hum Genet* 1999;105:442-3.
  44. Salem AH, Batzer MA. Distribution of the HIV resistance CCR5-Delta32 allele among Egyptians and Syrians. *Mutat Res* 2007;616:175-80.

**Cite this article as:** Chavhan AB, Pawar SS, Jadhao RG, Patil KG. Distribution of CC-chemokine receptor-5- $\Delta$ 32 allele among the tribal and caste population of Vidarbha region of Maharashtra state. *Indian J Hum Genet* 2013;19:65-70.

**Source of Support:** Nil, **Conflict of Interest:** None declared.